

Reducing Livestock Effects on Public Lands in the Western United States as the Climate Changes: A Reply to Svejcar et al

Robert L. Beschta · Debra L. Donahue · Dominick A. DellaSala ·
Jonathan J. Rhodes · James R. Karr · Mary H. O'Brien ·
Thomas L. Fleischner · Cindy Deacon Williams

Received: 26 January 2014 / Accepted: 15 March 2014
© Springer Science+Business Media New York 2014

Abstract Svejcar et al. (Environ Manage, 2014) offered several perspectives regarding Beschta et al. (Environ Manage 51:474–491, 2013)—a publication that addressed the interacting ecological effects of climate change and domestic, wild, and feral ungulates on public lands in the western United States (US)—by largely focusing on three livestock grazing issues: (1) legacy versus current day impacts; (2) grazing as a fire reduction tool; and (3) the complexity of grazing. Regarding these issues, we indicate that (1) legacy effects to western ecosystems were indeed significant and contemporary livestock use on public lands generally maintains or exacerbates many of those effects; (2) livestock grazing has been a major factor affecting fire frequency, fire severity, and ecosystem trajectories in the western US for over a century; and (3) the removal or reduction of grazing impacts in these altered ecosystems is the most effective means of initiating ecological recovery.

Svejcar et al. 2014 (see Literature cited).

R. L. Beschta (✉)
Department of Forest Ecosystems and Society, Oregon State
University, Corvallis, OR 97331, USA
e-mail: Robert.Beschta@oregonstate.edu

D. L. Donahue
College of Law, University of Wyoming, Dept. 3035,
1000 E. University Avenue, Laramie, WY 82071, USA

D. A. DellaSala
Geos Institute, 84 Fourth St., Ashland, OR 97520, USA

J. J. Rhodes
Planeto Azul Hydrology, P.O. Box 15286, Portland,
OR 97293, USA

J. R. Karr
102 Galaxy View Court, Sequim, WA 98382, USA

Svejcar et al. (Environ Manage, 2014) offer no evidence that livestock use is consistent with the timely recovery of grazing-degraded uplands, riparian areas, or stream systems. We thus conclude that public-land ecosystems can best persist or cope with a changing climate by significantly reducing ungulate grazing and related impacts.

Keywords Ungulates · Livestock grazing · Climate change · Public lands · Biodiversity · Restoration

Beschta et al. (2013) synthesized the ecological effects of climate change and ungulate grazing on western public lands, grounding their recommendations in ecological considerations and federal agency legal authority and obligations. Svejcar et al. (2014) suggest that Beschta et al. (2013) neither “present a balanced synthesis of the scientific literature” nor “reflect the complexities associated with herbivore grazing.” Svejcar et al. (2014) “dispute the notion that eliminating [livestock] grazing will provide a

M. H. O'Brien
Grand Canyon Trust, HC 64 Box 2604, Castle Valley,
UT 84532, USA

T. L. Fleischner
Environmental Studies and Natural History Institute, Prescott
College, 220 Grove Avenue, Prescott, AZ 86301, USA

C. D. Williams
Environmental Consultants, 4393 Pioneer Road, Medford,
OR 97501, USA

solution to problems created by climate change,” although we made no such claim. Instead, Beschta et al. (2013: p. 474) indicate that removal or reduction of livestock across large areas of public land will reduce a pervasive ecological stress, diminishing cumulative impacts on these ecosystems under climate change. We respond to three livestock grazing issues raised by Svejcar et al. (2014): (1) legacy versus contemporary effects, (2) fuels reduction and fire effects, and (3) grazing complexity and restoration.

Legacy and Contemporary Livestock Use have Caused Combined Effects

Livestock effects began soon after their introduction to semi-arid ecosystems west of the Rockies, which had evolved in an absence of large herds of ungulates (Mack and Thompson 1982). Contemporary grazing impacts (as described in Beschta et al. 2013) compound “legacy” effects, including altered fire regimes; biological soil crust loss, soil loss, and compaction; altered composition, structure, and function of upland, riparian, and stream biological communities; altered streamflow regimes; and reduced food-web support and physical habitat for terrestrial and aquatic biota (Blackburn 1984; Belsky et al. 1999; Kauffman and Pyke 2001; Belnap and Lange 2003; Fleischner 2010). Combined legacy and current grazing effects have left many streams with degraded riparian vegetation, accelerated bank erosion, widened and/or incised stream channels, and altered water quality (increased temperatures and sediment loads). These changes have many negative biological effects, including those on imperiled resident and anadromous fish (NRC 1996, 2002). Because the legacy effects of livestock were significant and extensive, contemporary grazing studies tend to *underestimate* ecological impacts, as they compare changes within already diminished systems (Fleischner 1994).

While some livestock impacts (e.g., soil loss or channel incision) may not be fully reversible in short timeframes, recovery of native plant communities and soil functions, which underpin terrestrial ecosystems, often occurs when the causes of degradation are removed or reduced. Despite changes in public land grazing practices over time, evidence indicates that contemporary livestock use thwarts ecological recovery. Cessation of livestock grazing can result in recovery of soil properties (Kauffman et al. 2004), riparian vegetation (Hough-Snee et al. 2013 and Fig. 1), and channel morphology (Herbst et al. 2012 and Fig. 1), relative to areas that continue to be grazed.

Riparian and stream ecosystems (Belsky et al. 1999; NRC 2002) and aspen (*Populus tremuloides*) communities (Seager et al. 2013) are biologically diverse and especially susceptible to the effects of livestock use. For



Fig. 1 A photopoint demonstrating vegetation and stream channel change following removal of livestock in the Northern Great Basin (Barnhardi Meadows, Hart Mountain National Antelope Refuge, Oregon). *Upper photo* was taken October, 1990 after approximately one century of livestock grazing during which livestock use was managed by the US Fish and Wildlife Service from 1940 to 1990. *Lower photo* was taken August, 2013 following 22 years of rest from livestock grazing. In this ecosystem, the reestablishment of willows (*Salix* spp.) and other wetland obligate species, as well as increased aspen recruitment, has occurred. Previously eroding stream banks have stabilized and stream channels narrowed, since the removal of livestock on the refuge. Photo credits: (*upper*) Bill Pyle and (*lower*) Schyler Reis

example, recent studies in Wyoming (Hessl and Graumlich 2002), Nevada (Kay 2003), Montana (Kimble 2007), Oregon (Seager 2010), and Utah (Kay 2011) point to high levels of livestock herbivory over many decades, sometimes in combination with wild ungulate impacts, as a major factor inhibiting aspen growth from seedling/sprouts into saplings and trees. These long-term effects hamper the ability of this tree species to persist in many western ecosystems. Livestock grazing also has widespread effects on the frequency and distribution of native grasses, forbs, and shrubs, and native wildlife species dependent upon those plants [e.g., sage-grouse (*Centrocercus urophasianus*); Manier et al. 2013].

Livestock Grazing is Not a Viable Tool for Reducing Fuels and Wildfire Effects

Livestock grazing in western US landscapes altered natural fire regimes by decreasing the frequency of low-severity fires beginning in the early 1900s (Swetnam and Betancourt 1998), making large areas prone to invasion by woody species and, in turn, more susceptible to high-severity fires (Chambers and Pellant 2008). Furthermore, cheatgrass (*Bromus tectorum*), an annual exotic, spreads rapidly throughout the Intermountain West as a result of livestock movement and overgrazing (Mack 1986), contributing to more frequent burning. Cheatgrass dominates nearly 70,000 km² in the Great Basin and is a component on an additional 250,000 km² (Diamond et al. 2012). Reisner et al. (2013) found that livestock grazing increases cheatgrass dominance in sagebrush steppe, livestock grazing is not likely a viable tool for reducing cheatgrass dominance because it promotes cheatgrass invasion, and reduced grazing may be one of the most effective means of conserving and restoring imperiled sagebrush ecosystems.

Although Livestock Grazing has Complex Ecological Consequences, Large-Scale Reductions in Grazing Effects are Likely to Reduce Cumulative Ecosystem Degradation

Recognizing the complexity of grazing issues was central to the synthesis and recommendations included in Beschta et al. (2013). Our analyses provided an integrative view of that complexity: we discussed three classes of ungulates (domestic, feral, wild), drawing examples from diverse vegetation types (shrub steppe, desert, conifer forest), and ecological attributes (such as water quality, hydrology, riparian areas, soils, hydrology, and biodiversity). Nevertheless, compelling reasons exist to single out livestock as a cause of ecological harm to native plant communities, terrestrial and aquatic habitats, and watershed processes (Belsky et al. 1999; Kauffman and Pyke 2001; Belnap and Lange 2003; NRC 2002). Livestock use is a principal cause of desertification in arid and semi-arid landscapes (Swetnam and Betancourt 1998; Belnap and Lange 2003; Fleischner 2010). It has the most extensive land-use footprint on western public lands (Beschta et al. 2013), and it continues at major public expense (Vincent 2012). Livestock production also contributes directly and indirectly to greenhouse gases, raising increasing concern about its climate effects (Ripple et al. 2014). The cessation or removal of factors that cause degradation or prevent recovery is the most effective and robust approach to ecological restoration (Kauffman et al. 1997). Unlike many stressors, livestock use is subject to human control.

Svejcar et al. (2014) assert that position statements by the American Fisheries Society (Armour et al. 1991) and the Wildlife Society (2010) “do not advocate removing livestock from western rangelands.” These position statements, however, as well as those of the Society for Conservation Biology (Fleischner et al. 1994), conclude that public-land grazing impacts need to be dramatically reduced to allow recovery of degraded ecosystems—an explicit recommendation of Beschta et al. (2013). Moreover, these position statements were developed without consideration of climate change effects.

Livestock use of public lands in the West remains a major stressor with effects of increasing concern under the overarching stressor of climate change. Its removal or reduction is an ecologically efficient and unambiguous approach for restoring resilience to large areas of these lands (see synthesis in Beschta et al. 2013). Because livestock grazing has diminished biodiversity and degraded ecosystems, the burden of proof for maintaining the grazing status quo is on Svejcar et al. (2014). But they offer no evidence that livestock use is compatible with the recovery of livestock-degraded uplands, riparian areas, or stream systems, or with retention of native species in arid and semi-arid ecosystems. Absent such evidence, and in the context of a changing climate, the only rational, effective, and direct alternative for ecologically restoring many western public lands is to reduce the effects of their most prominent stressor—livestock.

References

- Armour CL, Duff DA, Elmore W (1991) The effects of livestock grazing on riparian and stream ecosystems. *Am Fish Soc* 1991:7–11
- Belnap J, Lange OL (eds) (2003) *Biological soil crusts: structure, function, and management*. Springer, New York
- Belsky AJ, Matzke A, Uselman S (1999) Survey of livestock influences on stream and riparian ecosystems in the western United States. *J Soil Water Conserv* 54:419–431
- Beschta RL, Donahue DL, DellaSala DA, Rhodes JJ, Karr JR, O'Brien MH, Fleischner TL, Deacon Williams C (2013) Adapting to climate change on western public lands: addressing the ecological effects of domestic, wild, and feral ungulates. *Environ Manage* 51:474–491
- Blackburn WH (1984) Impacts of grazing intensity and specialized grazing systems on watershed characteristics and responses. In: Committee on developing strategies for rangeland management. *Developing strategies for rangeland management*. National Research Council, Westview Press, Boulder, Colorado, pp 927–983
- Chambers JC, Pellant M (2008) Climate change impacts on north-western and intermountain United States rangelands. *Rangelands* 30:29–33
- Diamond JM, Call CA, Devoe N (2012) Effects of targeted grazing and prescribed burning on community and seed dynamics of a downy brome (*Bromus tectorum*) dominated landscape. *Invasive Plant Sci Manage* 5:259–269

- Fleischner TL (1994) Ecological costs of livestock grazing in western North America. *Conserv Biol* 8:629–644
- Fleischner TL (2010) Livestock grazing and wildlife conservation in the American West: historical, policy and conservation biology perspectives. In: du Toit JT, Kock R, Deutsch JC (eds) *Wild rangelands: conserving wildlife while maintaining livestock in semi-arid ecosystems*. Blackwell Publishing, Boston, MA, pp 235–265
- Fleischner TL, Brown DE, Cooperrider AY, Kessler WB, Painter EL (1994) Society for conservation biology position statement: livestock grazing on public lands in the United States of America. *Soc Conserv Biol Newsl* 1:2–3
- Herbst DB, Bogan MT, Roll SK, Safford HD (2012) Effects of livestock exclusion on in-stream habitat and benthic invertebrate assemblages in montane streams. *Freshwater Biol* 57:204–217
- Hessl AE, Graumlich LJ (2002) Interactive effects of human activities, herbivory and fire on quaking aspen (*Populus tremuloides*) age structures in western Wyoming. *J Biogeogr* 29:889–902
- Hough-Snee N, Roper BB, Wheaton JM, Budy P, Lokteff RL (2013) Riparian vegetation communities change rapidly following passive restoration at a northern Utah stream. *Ecol Eng* 58:371–377
- Kauffman JB, Pyke DA (2001) Range ecology, global livestock influences. In: Levin SA (ed) *Encyclopedia of biological diversity*, vol 5. Academic Press, San Diego, CA, pp 33–52
- Kauffman JB, Beschta RL, Otting N, Lytjen D (1997) An ecological perspective of riparian and stream restoration in the Western United States. *Fisheries* 22:12–24
- Kauffman JB, Thorpe AS, Brookshire J, Ellingson L (2004) Livestock exclusion and belowground ecosystem responses in riparian meadows of eastern Oregon. *Ecol Appl* 14:1671–1679
- Kay CE (2003) Aspen management guidelines for BLM lands in north-central Nevada. Wildlife Management Services, Providence, UT
- Kay CE (2011) Why is aspen declining on Cedar Mountain? Measurement of aspen exclosures in southern Utah. Final Report, Utah State University Extension, Cedar City, UT
- Kimble DS (2007) Quaking aspen ecology on forest service lands north of Yellowstone National Park. MS thesis, Montana State University, Bozeman, MT
- Mack RN (1986) Alien plant invasion into the intermountain west: a case history. In: Mooney HA, Drake JA (eds) *Ecology of biological invasions of North America and Hawaii*. Springer, New York, pp 191–213
- Mack RN, Thompson JN (1982) Evolution in steppe with few large, hooved mammals. *Am Nat* 119:757–773
- Manier DJ, Wood DJA, Bowen ZH, Donovan RM, Holloran MJ, Juliusson LM, Manye KS, Oyler-McCance SJ, Quamen FR, Saher DM, Titolo AJ (2013) Summary of science, activities, programs, and policies that influence the rangewide conservation of greater sage-grouse (*Centrocercus urophasianus*). USDI, Geological Survey, Open File Report 2013–1098
- NRC (National Research Council) (1996) *Upstream: salmon and society in the Pacific Northwest*. National Academy Press, Washington, DC
- NRC (National Research Council) (2002) *Riparian areas: functions and strategies for management*. National Academy Press, Washington, DC
- Reisner MD, Grace JB, Pyke DA, Doescher PS (2013) Conditions favouring *Bromus tectorum* dominance of endangered sagebrush steppe ecosystems. *J Appl Ecol*. doi:10.1111/1365-2664.12097
- Ripple WJ, Smith P, Haberl H, Montzka SA, McAlpine C, Douglas H, Boucher DH (2014) Commentary: ruminants, climate change and climate policy. *Nat Clim Change* 4:2–5
- Seager ST (2010) Quaking aspen persistence in three Oregon landscapes. MS Thesis, Oregon State University, Corvallis, OR
- Seager ST, Cristina Eisenberg C, Clair SB (2013) Patterns and consequences of ungulate herbivory on aspen in western North America. *For Ecol Manage* 299:81–90
- Svejcar T, Boyd C, Davies K, Madsen M, Bates J, Sheley R, Marlow C, Bohnert D, Borman M, Mata-González R, Buckhouse J, Stringham T, Perryman B, Swanson S, Tate K, George M, Ruyle G, Roundy B, Call C, Jensen K, Launchbaugh K, Gearhart A, Vermeire L, Tanaka J, Derner J, Frasier G, Havsta K (2014) Western land managers will need all available tools for adapting to climate change, including grazing: a critique of Beschta et al. *Environ Manage*. doi:10.1007/s00267-013-0218-2
- Swetnam TW, Betancourt JL (1998) Mesoscale disturbance and ecological response to decadal climatic variability in the American southwest. *J Clim* 11:3128–3147
- Vincent CH (2012) *Grazing fees: overview and issues*. Congressional research service RS21232, Washington, DC
- Wildlife Society (2010) *Final position statement: livestock grazing on rangelands in the western US*. The Wildlife Society, Bethesda, MD